



UDP-GlcDH rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A23175	Rabbit	1 mg/ml	55 kD

Applications	WB,IHC,ELISA
Reactivity	Human,Mouse,Rat
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
Storage	-20°C/1 year
Specificity	UDG Polyclonal Antibody detects endogenous levels of UDG protein.
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Immunogen	The antiserum was produced against synthesized peptide derived from human UNG. AA range:191-240
Uniprot No	P13051
Alternative names	UNG; DGU; UNG1; UNG15; Uracil-DNA glycosylase; UDG
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Clonality	Polyclonal
Isotype	
Conjugation	
Background	uracil DNA glycosylase(UNG) Homo sapiens This gene encodes one of several uracil-DNA glycosylases. One important function of uracil-DNA glycosylases is to prevent mutagenesis by eliminating uracil from DNA molecules by cleaving the N-glycosylic bon
Other	Gene_name: UNG ; Protein_name: Uracil-DNA glycosylase; Expression: Epithelium,Hippocampus,Placenta,Uterus,

Product Images

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:



H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.