



CAMLG rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A11570	Rabbit	1 mg/ml	32 kD
Applications	WB,ELISA		
Reactivity	Human		
Dilution	WB 1:500-2000 ELISA 1:5000-20000		
Storage	-20°C/1 year		
Specificity	CAMLG Polyclonal Antibody detects endogenous levels of protein.		
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.		
Immunogen	Synthesized peptide derived from part region of human protein		
Uniprot No	P49069		
Alternative names			
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.		
Clonality	Polyclonal		
Isotype			
Conjugation			
Background	calcium modulating ligand(CAMLG) Homo sapiens The immunosuppressant drug cyclosporin A blocks a calcium-dependent signal from the T-cell receptor (TCR) that normally leads to T-cell activation. When bound to cyclophilin B, cyclosporin A binds an		
Other	Gene_name: CAMLG CAML ; Protein_name: Calcium signal-modulating cyclophilin ligand (CAML); Expression: Brain,Colon adenocarcinoma,		
Product Images			

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:

H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster



X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.