



DDB2 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A13481	Rabbit	1 mg/ml	46 kD

Applications	WB,ELISA
Reactivity	Human,Mouse
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage	-20°C/1 year
Specificity	DDB2 Polyclonal Antibody detects endogenous levels of protein.
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Immunogen	Synthesized peptide derived from human protein . at AA range: 260-340
Uniprot No	Q92466
Alternative names	
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Clonality	Polyclonal
Isotype	
Conjugation	
Background	damage specific DNA binding protein 2(DDB2) Homo sapiens This gene encodes a protein that is necessary for the repair of ultraviolet light-damaged DNA. This protein is the smaller subunit of a heterodimeric protein complex that participates in nucl
Other	Gene_name: DDB2 ; Protein_name: DNA damage-binding protein 2 (DDB p48 subunit) (DDBb) (Damage-specific DNA-binding protein 2) (UV-damaged DNA-binding protein 2) (UV-DDB 2); Expression: Brain,Epidermis,Epithelium,

Product Images

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:



H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.