



IDH3G rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A16040	Rabbit	1 mg/ml	43 kD

Applications	WB,ELISA
Reactivity	Human
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage	-20°C/1 year
Specificity	IDH3G Polyclonal Antibody detects endogenous levels of protein.
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Immunogen	Synthesized peptide derived from part region of human protein
Uniprot No	P51553
Alternative names	
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Clonality	Polyclonal
Isotype	
Conjugation	
Background	isocitrate dehydrogenase 3 (NAD(+)) gamma(IDH3G) Homo sapiens Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the
Other	Gene_name: IDH3G ; Protein_name: Isocitrate dehydrogenase [NAD] subunit gamma, mitochondrial (EC 1.1.1.41) (Isocitric dehydrogenase subunit gamma) (NAD(+)-specific ICDH subunit gamma); Expression: Brain,Heart,Placenta,Skin,Soares liver spleen 1NFLS,

Product Images

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:



H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.