



HCAR2 rabbit pAb antibody

Catalog No :	Source:	Concentration :	Mol.Wt. (kD):
A15511	Rabbit	1 mg/ml	39 kD

Applications	WB,ELISA
Reactivity	Human
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage	-20°C/1 year
Specificity	HCAR2 Polyclonal Antibody detects endogenous levels of protein.
Source / Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Immunogen	Synthesized peptide derived from human protein . at AA range: 260-340
Uniprot No	Q8TDS4
Alternative names	
Form	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Clonality	Polyclonal
Isotype	
Conjugation	
Background	developmental stage:Expression in neutrophils occurs in the late terminal differentiation phase.,function:Acts as a high affinity receptor for both nicotinic acid (also known as niacin) and (D)-beta-hydroxybutyrate and mediates increased adiponectin secre
Other	Gene_name: HCAR2 GPR109A HCA2 HM74A NIACR1 ; Protein_name: Hydroxycarboxylic acid receptor 2 (G-protein coupled receptor 109A) (G-protein coupled receptor HM74A) (Niacin receptor 1) (Nicotinic acid receptor); Expression: Pancreas,

Product Images

Application Key:

W-Western IP-Immunoprecipitation IHC-Immunohistochemistry ChIP-Chromatin Immunoprecipitation
IF-Immunofluorescence F-Flow Cytometry E-P-ELISA-Peptide

Species Cross-Reactivity Key:



H-Human M-Mouse R-Rat Hm-Hamster Mk-Monkey Vir-Virus Mi-Mink C-Chicken Dm-D. melanogaster
X-Xenopus Z-Zebrafish B-Bovine Dg-Dog Pg-Pig Sc-S. cerevisiae Ce-C. elegans Hr-Horse All-All
Species Expected

Trademarks

All product names and trademarks are the property of their respective owners.

Regulatory Disclaimer

For life science research only. Not for use in diagnostic procedures.

Contact and Support:

To ask questions, solve problems, suggest enhancements and report new applications, please visit our [Online Technical Support Site](#).

To call, write, fax, or email us, please visit www.aabsci.com, contact information will be displayed.